



# Robustness checks in PLS-SEM: A review of recent practices and recommendations for future applications in business research

Santha Vaithilingam<sup>a,\*</sup>, Chu Sun Ong<sup>b</sup>, Ovidiu I. Moisescu<sup>c</sup>, Mahendhiran S. Nair<sup>a</sup>

<sup>a</sup> Sunway Institute for Global Strategy & Competitiveness & Sunway Business School, Sunway University, Malaysia

<sup>b</sup> Department of Innovation Science, School of Environment and Society, Tokyo Institute of Technology, Japan

<sup>c</sup> Department of Marketing, Faculty of Economics and Business Administration, Babeş-Bolyai University, Romania

## ARTICLE INFO

### Keywords:

PLS-SEM  
Nonnormality  
Endogeneity  
Unobserved heterogeneity  
Nonlinearity  
Heteroskedasticity

## ABSTRACT

For the last decade, scholars have employed partial least squares structural equation modeling (PLS-SEM) extensively in business research. However, when applying PLS-SEM, researchers need to perform various robustness checks before and after model estimation. This study showcases the findings of a review of PLS-SEM use in business research, by examining papers published between 2016 and 2021 in business journals. The study explores the extent to which researchers have performed robustness checks regarding nonnormality, endogeneity, unobserved heterogeneity, nonlinearity, and heteroskedasticity. The findings highlight that statistical rigor remains a serious problem in business-related studies employing PLS-SEM. Despite some encouraging improvements in the last few years, the vast majority of recent business-related studies using PLS-SEM have systematically overlooked robustness checks. This study calls for continued emphasis on the importance of robustness checks and the correct application of appropriate techniques, providing recommendations and guidelines for future PLS-SEM applications.

## 1. Introduction

For the last decade, partial least squares structural equation modeling (PLS-SEM) has risen in popularity as a method of choice for investigating intricate relationships between observed and latent constructs in social science research. This increased popularity is evidenced by the exponential growth in the number of PLS-SEM applications published in major social science journals (Hair et al., 2022). Within the domain, PLS-SEM is particularly appealing and it is employed extensively in business research in a wide range of subjects, such as marketing (Sarstedt et al., 2022a), international management (Richter et al., 2022), strategic management (Hair et al., 2012), supply chain management (Kaufmann & Gaeckler, 2015), operations management (Bayonne et al., 2020), human resource management (Ringle et al., 2020), family business management (Hair et al., 2021), management information systems (Hair et al., 2017), knowledge management (Cepeda-Carrión et al., 2018), technology adoption and use (Vaithilingam et al., 2022) and hospitality management (Ali et al., 2018).

Compared to other SEM methods that used to be more prominent in the past, particularly covariance-based structural equation modeling (CB-SEM), PLS-SEM provides several advantages to researchers. It

facilitates the estimation of intricate models using relatively little data, with no imposed distributional assumptions, this being especially beneficial for business researchers who often rely on relatively small samples and nonnormal data (Hair et al., 2019b). In addition, PLS-SEM enables researchers to handle both reflectively and formatively specified measurements in their estimated models to test hypothesized relationships and to emphasize prediction, all at the same time (Cepeda-Carrión et al., 2016). This is of particular interest in business studies in which confirmation and explanation, as typical requisites of academic research, need to be accompanied by prediction in order to derive managerial implications (Henseler, 2018; Shmueli et al., 2016). Moreover, the availability of intuitive and user-friendly software packages (Sarstedt & Cheah, 2019), along with various papers and textbooks providing guidelines for applying the technique (e.g., Hair et al., 2022), has made PLS-SEM even more appealing to social science researchers, in general, and business researchers, in particular.

However, when applying PLS-SEM, researchers need to perform various robustness checks before and after model estimation, as there are several instances, such as nonnormality, endogeneity, unobserved heterogeneity, nonlinearity, and heteroskedasticity, that might threaten the validity of the results. Thus, even though PLS-SEM is a

\* Corresponding author at: No. 5, Jalan Universiti, Bandar Sunway 47500, Selangor Darul Ehsan, Malaysia.

E-mail address: [santhaav@sunway.edu.my](mailto:santhaav@sunway.edu.my) (S. Vaithilingam).

nonparametric method and does not require normally distributed data, extremely nonnormal data can lead to misleading results regarding the model parameters' statistical significance (Hair et al., 2022). Furthermore, endogeneity can lead to inaccurate path coefficients in PLS-SEM models. This is particularly problematic when testing hypothesized relationships, especially in studies focusing on confirmation and explanation (Hult et al., 2018; Papies et al., 2017). In addition, assuming that the investigated population is homogeneous, which is often unrealistic in business research, can lead to incorrect conclusions if a heterogeneous sample is analyzed on the aggregate level (Sarstedt et al., 2022b). Moreover, overlooking nonlinearities in the structural model when they can better depict the relationships between constructs, and erroneously assuming linearity for all paths in the PLS-SEM model, can lead to underestimating the true relationships or even wrongly flagging them as non-significant (Basco et al., 2022). Ultimately, although the method's nonparametric nature does not imply meeting rigorous homoskedasticity assumptions, extremely heteroskedastic data should be handled carefully, as PLS-SEM is a regression-based method.

Along with the rise of PLS-SEM as a method of choice, research has produced various additional methods for evaluating the robustness of PLS-SEM output. Nevertheless, despite their extensive documentation in the literature, business research using PLS-SEM has been slow in adopting those robustness checks. For instance, Latan (2018) and Sarstedt et al. (2020) highlighted the very limited use of robustness checks in hospitality and tourism research, providing guidelines to check for endogeneity, unobserved heterogeneity, and nonlinearity in a PLS-SEM framework.

Considering the utmost importance of robustness checks in PLS-SEM, this paper showcases the findings of a review of PLS-SEM use in business research, by examining papers published between 2016 and 2021 in high-quality journals. Our analysis examines the extent to which business research has incorporated robustness checks (i.e., concerning nonnormality, endogeneity, unobserved heterogeneity, nonlinearity, and heteroskedasticity) in the application of PLS-SEM, and provides recommendations and guidelines for conducting such checks before and after model estimation in future PLS-SEM studies. To achieve this objective, we attempt to answer the following questions: (i) What type of robustness checks were performed, (ii) which techniques were used, and (iii) is there a trend in performing and reporting results of robustness checks.

While we acknowledge the importance of other robustness checks (e.g., necessary condition analysis, common method variance, confirmatory tetrad analysis, model comparisons), our selection is the result of deliberate prioritization based on the relevance of such checks within the conventional regression analysis framework, considering that PLS-SEM is a regression-based method. Practical considerations related to time constraints and data availability also played a role in the decision to highlight these selected checks.

In our investigation, the robustness checks are categorized according to the point of time at which the analysis is undertaken. Thus, normality is assessed before model estimation. This is to ensure that extremely nonnormal data does not lead to misleading results, and also to ascertain the suitability of PLS-SEM due to its robustness when analyzing non-normal data. On the other hand, endogeneity, unobserved heterogeneity, nonlinearity, and heteroskedasticity are evaluated after model estimation to warrant the validity of the results.

The paper is structured as follows. We start by describing the protocols of the systematic literature review (SLR). We subsequently outline our SLR findings, depict the prevalence of robustness checks in the investigated papers, and discuss the techniques used for each robustness check. Finally, we provide recommendations for future PLS-SEM applications in business research.

## 2. Methodology

We conducted an SLR, following the guidelines of Kitchenham et al.

(2009) who pointed out that the SLR functions as an important tool for assessing the methodological quality of research. Given the plethora of empirical studies that use PLS-SEM, and the guidelines published to help researchers use and report PLS-SEM results, the question of statistical rigor in those studies remains elusive (Hair et al., 2024; Hult et al., 2018; Sarstedt et al., 2020).

Since the scope of the review included all papers that employed the PLS-SEM methodology, the query had to contain all the different terminologies used to refer to PLS-SEM. Hence, to obtain a comprehensive list of results, the following search strings were used: "(PLS-SEM) OR (PLS) OR (PARTIAL LEAST SQUARES)." The search was conducted within Clarivate Analytics' Web of Science Core Collection, a database that covers more than 21,100 leading journals.

Several restrictions were applied to filter irrelevant results while maintaining the reliability of the SLR results. The initial search criteria were set to locate papers that are published (i) in English, (ii) in academic journals, (iii) in a business field of study, and (iv) for the past six years (2016–2021). Further criteria of inclusion applied to the search results are as follows:

(vi) The paper was published in a high-ranked journal, namely a Q1 journal as per the 2020 SCImago Journal Rank (SJR) indicator<sup>1</sup> (SCImago, n.d.), as methodological rigor characterizes high-quality journals (West & Rich, 2012); (vii) the paper was empirical in nature (only papers with applications of PLS-SEM were considered; theoretical, conceptual, or "guidelines" papers were excluded).

The initial search returned 2,725 articles. After the exclusion of articles based on the established selection criteria, a total of 1,228 articles remained. These articles were published in 76 unique business journals. Appendix A shows these journals, with the reported instances of robustness checks. The flow chart highlighting the different phases of our SLR is shown in Fig. 1.

To enhance comparability and enable the examination of the progression in robustness checks practices within the business literature, we subsequently partitioned the corpus of articles into two distinct sub-periods, each comprising an equivalent span of years (i.e., 2016–2018 and 2019–2021).

## 3. Findings

Of the 1,228 papers selected, we found that 182 (14.9 %) reported at least one of the diagnostic checks within the scope of our review. In these 182 papers, a total of 245 instances of robustness checks were reported. In the next sections, we discuss our findings and their key implications for each category of robustness checks.

### 3.1. Normality

Researchers use bootstrapping for hypothesis testing in PLS-SEM (Hair et al., 2022), rendering PLS-SEM a distribution-free technique. Unlike CB-SEM, PLS-SEM does not require normally distributed data which is particularly appealing for business researchers who often rely on nonnormal data (Hair et al., 2019b). However, when using PLS-SEM, extremely nonnormal data can lead to misleading results regarding the statistical significance of the parameters (Guenther et al., 2023; Hair et al., 2022). For this reason, and also to support their method of choice, researchers employing PLS-SEM should assess the normality of their data. Nevertheless, motivating the choice of PLS-SEM over alternative methods primarily on the basis of nonnormality alone is not sufficient (Sarstedt et al., 2022a). Additional robust arguments may encompass considerations such as the meaningfulness of indicator residual variances for the constructs in the model (Guenther et al., 2023), or the

<sup>1</sup> Our classifications of "high quality" journals are based solely on the SJR score. Therefore, they may potentially exclude journals considered high quality by other ranking systems.




Fig. 1. Flow diagram.

predictive nature of the research objective (Hair et al., 2019c; Ringle et al., 2023).

Normality tests can be categorized according to the type of normality being tested, namely univariate or multivariate normality. Univariate normality is a requisite for multivariate normality (DeCarlo, 1997). Consequently, potential users of PLS-SEM should first assess univariate normality, checking whether individual indicators or items are normally distributed. If univariate normality is not rejected, users can then proceed to evaluate the stronger assumption of multivariate normality.

There are various approaches for univariate normality assessment, ranging from graphical methods to summary measures, and statistical tests. However, graphical methods including the Q-Q plot and P-P plot, have been criticized for the subjectivity of their interpretation (Loy et al., 2016). The common statistical tests that can be employed to assess normality are the Shapiro-Wilk and Kolmogorov-Smirnov tests. The former test has a higher power and is therefore recommended for small samples (Razali & Wah, 2011). Nevertheless, as these tests cannot indicate the extent of departure from normality, PLS-SEM users should examine skewness and kurtosis values. Skewness and kurtosis are summary statistics that measure the extent of departures from normality and can be used in both a descriptive and an inferential manner. Values of

skewness and kurtosis outside the range of  $-1$  to  $1$  denote significant deviation from normality. However, considering the robust performance of PLS-SEM when data are nonnormal, values between  $-2$  and  $+2$  can generally be considered acceptable (Hair et al., 2022).

As for multivariate normality assessment, researchers can employ tests, such as those developed by Mardia (1970, 1974), Small (1980), or Srivastava (1984). However, this assessment is not required if univariate normality, a prerequisite for multivariate normality, is rejected.

Our review of studies using PLS-SEM in business research shows that only 129 of 1,228 studies (10.5 %) assessed the normality of their data. Moreover, this situation has not improved in more recent years, with 90 of 858 studies (10.5 %) published between 2019 and 2021 performing such checks, compared to 39 out of 370 papers (10.5 %) published between 2016 and 2018.

As Table 1 shows, examining skewness and kurtosis values, as recommended by Hair et al. (2022), consistently proved to be the most used approach for normality assessment (54 of the 129 studies, 41.7 %). Statistical tests were also frequently performed to test for normality, the most common of which were the Shapiro-Wilk and Kolmogorov-Smirnov tests. Nevertheless, by not examining skewness and kurtosis values, more than half of the 129 studies failed to remove doubts associated with extreme nonnormalities that might have led to misleading results. Moreover, 23 of the 129 studies (17.8 %) explicitly claimed that normality checks were performed and that their data were nonnormal but provided no information on the type of assessment. In fact, studies that exploited the non-parametric nature of PLS-SEM to justify the use of the method, by making rhetoric statements (e.g., “PLS-SEM is chosen instead of CB-SEM, as PLS-SEM can handle nonnormally distributed data”) without conducting the appropriate tests or reporting results of the tests, proved to be numerous (236 out of the total of 1,228 reviewed studies).

### 3.2. Endogeneity

Endogeneity occurs when a predictor (explanatory) variable in a regression equation is correlated with the error term of the dependent variable (Wooldridge, 2013). If that is the case, it can lead to inaccurate path coefficients in PLS-SEM models (Papies et al., 2017), which is particularly problematic when testing hypothesized relationships. Even though endogeneity can have several roots, it is usually the result of omitted latent variables that correlate with both the dependent variable and a predictor variable in the PLS-SEM model (Sarstedt et al., 2020). One of the approaches to address this is to employ control and instrumental variables (Ebbes et al., 2016). However, such variables are often unavailable or difficult to find and, consequently, PLS-SEM users can employ the more accessible Gaussian copula approach, which enables PLS-SEM users to assess the correlation between explanatory or predictor constructs and the error terms of dependent ones (Hair et al.,

Table 1  
Number of instances normality checks were performed.

Technique used	2016–2018		2019–2021	
	Count	%	Count	%
Skewness and kurtosis	14	28	40	39
Details not provided	9	18	14	14
Kolmogorov-Smirnov	9	18	14	14
Shapiro-Wilk	7	14	11	11
Mardia	5	10	11	11
Graphical method	0	0	6	6
Jarque-Bera	0	0	4	4
Multivariate skewness and kurtosis	2	4	1	1
Kurtosis	0	0	2	2
Small	1	2	0	0
Srivastava	1	2	0	0
Others	2	4	0	0
<b>Total</b>	<b>50</b>	<b>100</b>	<b>103</b>	<b>100</b>

2024).

Our review of papers employing PLS-SEM in business research shows that endogeneity checks were performed in only 28 of 1,228 studies (2.3 %). However, from this perspective, PLS-SEM users' rigor has improved significantly in more recent years, since endogeneity checks were performed in 25 of 858 studies (2.9 %) published between 2019 and 2021, compared to only 3 of 370 (0.8 %) published between 2016 and 2018. This is an encouraging finding, considering that a previous review of studies published between 2008 and 2017 in several of the highest ranked marketing journals revealed that none of the investigated PLS-SEM applications had considered any endogeneity assessment (Hult et al., 2018).

As Table 2 shows, studies that checked for endogeneity employed various tests, such as the Gaussian copula approach (Park & Gupta, 2012), the Durbin-Wu-Hausman (DWH) test, the Heckman test, or the nonlinear bivariate causality direction ratio (NLBCDR) (Kock, 2015). Among the endogeneity tests identified in our SLR, the Gaussian copula approach proved to be the PLS-SEM users' preferred test, most probably due to its instrumental variable (IV)-free nature. The DWH test, on the other hand, is less popular, as it requires identification of an appropriate instrumental variable. Moreover, the DWH is a parametric test, while PLS-SEM is known for its non-parametric nature.

Kock's NLBCDR method is still experimental (Kock, 2022) and is currently useful for testing endogeneity arising from reverse causality. Besides simultaneity, which comprises reverse causality, other sources of endogeneity include omitted variable, measurement error, and selection bias (Wooldridge, 2010). Endogeneity caused by these sources might not be detected using Kock's NLBCDR approach. The use of the Heckman test, on the other hand, is limited to detecting endogeneity caused by sample selection bias (Certo et al., 2016; Wooldridge, 2013). Ramsey's Regression Equation Specification Error Test (RESET) has also been suggested for endogeneity assessment. This application is, however, inappropriate, as RESET is a general misspecification test for linear regression models (Ramsey, 1969) and is only useful for detecting functional form misspecification, particularly the existence of neglected nonlinearities between dependent and independent variables that exist in the regression equation (for details, see Wooldridge, 2010).

Some papers account for endogeneity at the research design phase to ensure that all possible factors that have a potential impact on the outcome (dependent) variable are considered in their conceptual model. While a theoretical treatment of endogeneity was acceptable in the past (Guide & Ketokivi, 2015), the use of statistical evidence such as that based on the IV-free Gaussian copula approach (Park & Gupta, 2012) has become desirable. While it is IV-free, able to handle multiple endogenous regressors, and implemented in PLS-SEM software packages (e.g., SmartPLS 4.0 (Ringle et al., 2022)), Becker et al. (2022) advise researchers to be careful in applying the Gaussian copula approach. The authors highlight that the approach misleads if the following three conditions are not fulfilled: (i) nonnormality of the endogenous regressor(s), (ii) normality of the error term, and (iii) the error term and the endogenous variable follow a Gaussian copula correlation structure. The Cramér-von Mises (also implemented in PLS-SEM software packages

such as SmartPLS 4.0 (Ringle et al., 2022)) and Anderson-Darling tests proved to have the best performance in testing for the nonnormality of the endogenous regressor (Becker et al., 2022). The normality of the error term condition can be tested by ensuring the regression residuals from the PLS-SEM structural model are normally distributed. However, the correlation structure required by the third condition cannot be observed whereby researchers must resort to theoretical considerations. Even when these three conditions can be justified, the Gaussian copula approach has a further minimum sample size requirement of 200 if skewness of the endogenous regressor is higher than 2; 1,000 if skewness is higher than 0.8; and 2,000 if skewness is below 0.8 (Becker et al., 2022).

Endogeneity assessment is not compulsory in all applications of PLS-SEM and is dependent on whether the objective of the study is confirmatory or predictive (Ebbes et al., 2011). In confirmatory-type studies, controlling for endogeneity is important to ensure the reliability of hypothesis testing results. In contrast, endogeneity is less of a concern in predictive-type studies where hypothesis testing is not the main objective of the research, and controlling for it may have an impact on the model's out-of-sample predictive power (Papies et al., 2017). Nonetheless, the line between confirmatory and predictive studies is often blurred in PLS applications (Schuberth et al., 2022). This calls for research to identify the approaches to handle endogeneity, by considering the interplay between the two research paradigms in the PLS-SEM context (Hair et al., 2019c). Nevertheless, it is still important for PLS-SEM users to determine, a priori, the prevailing intent of the study as either confirmatory or predictive, given the distinctive difference between the two types of studies in every stage of the statistical modeling process (Shmueli, 2010).

### 3.3. Unobserved heterogeneity

In PLS-SEM studies, assuming that the investigated population is homogeneous (which is often unrealistic in business research) can lead to incorrect conclusions if a heterogeneous sample is analyzed on the aggregate level (Sarstedt et al., 2022b). One way of addressing heterogeneity is to partition the data based on observable characteristics of the population and to estimate the model separately for each partition, thus accounting for observed heterogeneity. However, very often the cause of heterogeneity is unknown a priori and, consequently, unobserved heterogeneity needs to be accounted for. If such checks indicate no substantial repercussions affecting the results, the data can be analyzed on the aggregate level and their output can be generalized. However, if unobserved heterogeneity is detected, it needs to be treated accordingly.

In order to detect and handle unobserved heterogeneity appropriately, PLS-SEM users have a multitude of specific methods at their disposal, generally termed as latent class techniques. The most notable such technique is finite mixture PLS (FIMIX-PLS; Sarstedt et al., 2011). The method enables researchers to reveal the existence of heterogeneity reliably and to determine the number of segments correctly, using information criteria (e.g., jointly considering AIC3 and CAIC) and classification criteria (e.g., entropy statistic) (Sarstedt et al., 2022b). The technique's implementation in popular PLS-SEM software packages (e.g., SmartPLS), along with the many articles providing guidelines for applying it (e.g., Hair et al., 2016; Matthews et al., 2016), has made it readily available for PLS-SEM users for many years.

However, when heterogeneity is detected, identifying the actual segment structure correctly is an important limitation of FIMIX-PLS (Ringle et al., 2014; Sarstedt et al., 2022b), more so when formative constructs are involved (Becker et al., 2013). To overcome this limitation, research has proposed several additional techniques, such as the response-based procedure for detecting unit segments in PLS (REBUS-PLS, Esposito Vinzi et al., 2008), the prediction-oriented segmentation in PLS-SEM (PLS-POS, Becker et al., 2013), the PLS genetic algorithm segmentation (PLS-GAS, Ringle et al., 2014), and the PLS iterative reweighted regressions (PLS-IRRS, Schlittgen et al., 2016). However,

**Table 2**  
Number of instances endogeneity checks were performed.

Technique Used	2016–2018		2019–2021	
	Count	%	Count	%
Gaussian copula approach	0	0	10	38
Hausman/Durbin-Wu-Hausman	3	100	5	19
NLBCDR*	0	0	5	19
Heckman	0	0	2	8
Control variable approach	0	0	2	8
Ramsey*	0	0	2	8
<b>Total</b>	<b>3</b>	<b>100</b>	<b>26</b>	<b>100</b>

\* Inappropriate technique for the intended robustness check.

instead of employing any of these additional techniques exclusively, researchers are advised to adopt a tandem approach (Sarstedt et al., 2022b), using FIMIX-PLS to detect heterogeneity and determine the number of segments, and to further use its results as a starting point for applying additional latent class techniques (i.e., using PLS-POS, PLS-REBUS, PLS-GAS, or PLS-IRRS). Moreover, since simulation studies have shown that PLS-POS provides favorable outcomes compared to its alternative techniques, and considering its availability in popular PLS-SEM software packages (e.g., SmartPLS), PLS-SEM users are recommended to use FIMIX-PLS and PLS-POS in tandem when heterogeneity is detected (Hair et al., 2024). Based on PLS-POS results, it is then mandatory to conduct ex-post analyses in order to link the latent segment structure to observable characteristics in the data (Sarstedt et al., 2022b), using techniques such as contingency table analyses and exhaustive chi-squared automatic interaction detector (CHAID) (Ringle et al., 2010), fuzzy-set qualitative comparative analysis (fsQCA) (Mikalef & Pateli, 2017), or logistic regression analyses (Dessart et al., 2019).

Our review shows that only 30 of 1,228 studies (2.4 %) performed unobserved heterogeneity checks for their data. Compared to a recent review on PLS-SEM use in marketing research (Sarstedt et al., 2022a), which found that 4.18 % of studies conducted latent class analyses, our review within the wider field of business research reveals an even less rigorous stand on this issue. Moreover, this situation has worsened in more recent years, with only 20 of 858 studies (2.3 %) published between 2019 and 2021 performing such checks, compared to 10 of 370 papers (2.7 %) published between 2016 and 2018.

As seen in Table 3, FIMIX-PLS dominates among the methods used for uncovering unobserved heterogeneity, which is in line with previous methodological recommendations. However, according to our review, only 4 of the 11 studies that had used FIMIX-PLS and that had detected unobserved heterogeneity further employed PLS-POS (or other additional latent techniques) to better depict the actual segment structure. In addition, among the five studies employing PLS-POS, two have done so without adopting the recommended tandem approach (i.e., FIMIX-PLS, followed by PLS-POS).

Unfortunately, our review emphasizes that business research has been slow in responding to recent calls to check for unobserved heterogeneity (e.g., Sarstedt et al., 2020, first published online in 2019). This is despite the wide availability of latent class techniques, both considering software implementation, and guideline papers. Hopefully, more recent calls (e.g., Sarstedt et al., 2022b), along with our current strong recommendation to account for unobserved heterogeneity, will have a significant impact in the near future.

### 3.4. Nonlinearity

Despite many past studies assuming monotonic positive linear relationships between constructs, Pierce and Aguinis (2013) pointed out that such relationships can become asymptotic and, consequently, negative with the increase in the value of the independent variables, resulting in a curvilinear relationship that they refer to as the “too-much-of-a-good-thing effect” (TMGT effect). For a nonlinear

relationship between two variables, the effect size depends on the magnitude of change in the independent variable (Hair et al., 2024), that is, the changes in the slopes are dependent on the level of the variable (Basco et al., 2022). Thus, understanding the true underlying relationship between the independent and outcome variables is important, as the functional form will be dependent on the pattern of the relationship. Overlooking nonlinearities in the structural model when they can better depict the relationships between constructs, and erroneously assuming linearity for all paths in a PLS-SEM model can lead to underestimating the true relationships, or even wrongly flagging them as non-significant (Basco et al., 2022).

The RESET test (Ramsey, 1969) is the most prominent technique to be used for checking nonlinearities in PLS-SEM (Latan et al., 2018; Svensson et al., 2018). It involves estimating the PLS-SEM model and using the resulted latent scores as inputs for RESET, which can be implemented in standard software (e.g., SPSS, E-Views, or Stata; Sarstedt & Mooi, 2014). RESET tests are based on fitted values from a linear model. For this reason, the second stage is required to assess if a nonlinear specification for the relationships of the latent constructs in the structural model results in significant coefficients (Latan et al., 2018). This can be confirmed by adding the interaction terms in the structural model and reanalyzing the structural model based on bootstrapping (Hair et al., 2024). According to Hair et al. (2018), especially in social sciences, modeling quadratic effects should suffice, as other higher order polynomials of the independent constructs are very challenging.

Our review of articles applying PLS-SEM in business research shows that only 16 of 1,228 studies (1.3 %) accounted for nonlinearity in the investigated models, among which three studies did not actually report the results of their nonlinearity checks. There is, however, a slight improvement over time, with 12 of 858 studies (1.4 %) published between 2019 and 2021 performing such checks, compared to 4 of 370 papers (1.1 %) published between 2016 and 2018. Our findings are consistent with those of Sarstedt et al. (2020) who highlighted that most studies using PLS-SEM in hospitality and tourism research had not considered nonlinear effects, and also with those of Basco et al. (2022) who emphasized that family business research has only rarely considered nonlinear effects.

As Table 4 shows, the RESET technique for nonlinearity assessment has gained attention in more recent years, as all four studies employing the method were published between 2019 and 2021. However, only two of these studies have actually applied the recommended methodological rigor by completing both the RESET and quadratic effect assessments.

Just as in the case of unobserved heterogeneity, our review reveals once more that business research has been slow in responding to recent calls for robustness checks, including nonlinearity assessment. A possible explanation could be the cumbersome process of combining the use of several software packages (i.e., RESET), and the multi-staged nature of the full procedure (i.e., RESET followed by quadratic effect assessment). However, there is optimism that the recent calls for robustness checks (e.g., Sarstedt et al., 2022a), along with our strong recommendation to account for nonlinear effects, will give nonlinearity

**Table 3**  
Number of instances unobserved heterogeneity checks were performed.

Technique Used	2016–2018		2019–2021	
	Count	%	Count	%
FIMIX-PLS*	9	82	16	73
PLS-POS	2	18	3	14
REBUS-PLS	0	0	2	9
Details not provided	0	0	1	5
<b>Total</b>	<b>11</b>	<b>100</b>	<b>22</b>	<b>100</b>

\*Of the 25 studies that conducted FIMIX-PLS, 11 detected unobserved heterogeneity.

**Table 4**  
Number of instances nonlinearity checks were performed.

Technique Used	2016–2018		2019–2021	
	Count	%	Count	%
Details not provided	1	25	2	17
Graphical method	1	25	1	8
RESET	0	0	2	17
Quadratic effect assessment	1	25	1	8
RESET and quadratic effect assessment	0	0	2	17
ANOVA (SPSS)	0	0	2	17
Curve estimation	1	25	1	8
Warp3 algorithm	0	0	1	8
<b>Total</b>	<b>4</b>	<b>100</b>	<b>12</b>	<b>100</b>

checks a boost in future studies.

### 3.5. Heteroskedasticity

PLS-SEM users might be unfamiliar with the concept of heteroskedasticity (HSK), even though the issue of observed or unobserved heterogeneity has been fairly established in the PLS-SEM literature. Heterogeneity is about the estimated path coefficient and if these coefficients vary significantly across subgroups of the population. HSK, on the other hand, refers to whether the variability around the estimated path coefficient differs across the different values of the antecedents or predictor variables in the model. Simply put, heterogeneity addresses the first moment (i.e., the mean), while HSK addresses the second moment (i.e., the variance). Given this property, HSK is also known as heterogeneity in the variance.

To account for HSK, PLS-SEM users can employ tests, such as the Levene test and the Breusch-Pagan and Koenker test. The Levene test is the most used HSK test and is meant for testing the univariate form of HSK. The test involves comparing the variance of a metric variable across levels of a nonmetric variable. The Breusch-Pagan and Koenker test, on the other hand, is meant for testing homogeneity of variance between two metric variables, which is a commonly faced issue in regression analysis (Hair et al., 2019a).

Although relatively few papers checked for the assumption of constant variance in the error terms or homoskedasticity, the numbers of studies came close to those checking for nonlinearity. Our review shows that 12 of 1,228 studies (1 %) performed HSK tests. Nevertheless, the adoption of such checks has improved in more recent years, since nine of 858 studies (1.1 %) published between 2019 and 2021 performed such checks, compared to three of 370 papers (0.8 %) published between 2016–2018. Table 5 shows that, as expected, the Levene test has been by far the most widely used to account for HSK.

A review of commonly cited PLS-SEM textbooks (e.g., Hair et al., 2024; Hair et al., 2022) and guideline articles (e.g., Hair et al., 2019b; Sarstedt et al., 2022a) did not mention whether the violation of the HSK assumption would pose a serious threat to the validity of PLS-SEM results. The fact that PLS-SEM is regression based but at the same time utilizes the bootstrapping procedure in generating the test statistics, makes the relevance of the HSK assumption as stipulated in the Gauss–Markov theorem unclear. In the interests of the diverse PLS-SEM community, which consists of researchers from many disciplines other than statistics, future research should address this gap.

## 4. Conclusions

Our study highlights that statistical rigor remains a serious problem in business-related studies using PLS-SEM. Only one in ten such studies published between 2016 and 2021 assessed the normality of their data, while endogeneity, unobserved heterogeneity, nonlinearity, or HSK checks were performed in only 2.3 %, 2.4 %, 1.3 %, and 1 %, respectively, of the 1,228 reviewed papers. Despite a significant improvement in what concerns endogeneity checks, as well as a slight advancement regarding nonlinearity and HSK examinations, the vast majority of studies using PLS-SEM in business research have largely ignored

robustness checks.

In addition, even when robustness checks were conducted, some studies showed uncertainty not only regarding the techniques to be used, but also regarding their appropriate context. This is more concerning, as misapplying such techniques can result in misleading conclusions or failing to fully depict the relationships between constructs in PLS-SEM models.

When performing robustness checks, researchers are advised not to be selective or adopt a “cherry-picking” behavior, but to be rigorous and transparent in performing and reporting the results of the checks performed. Otherwise, the validity of their findings will be called into question.

In line with recent calls for robustness checks (e.g., Sarstedt et al., 2022a), our strong recommendation for business researchers using PLS-SEM is to assess data normality and to account for endogeneity, unobserved heterogeneity, and nonlinearity. Table 6 provides a summary of the recommended techniques for each type of robustness check.

As for HSK, our SLR revealed several instances of such checks being performed. However, to date, no studies have addressed the extent to which HSK can affect the validity of PLS-SEM results. Hence, considering that PLS-SEM is a regression-based method, future research should examine the consequences of using extremely heteroskedastic data when assessing models in the PLS-SEM framework.

Moving forward, in order to encourage rigor among business researchers using PLS-SEM, PLS-SEM software developers should strive to include built-in functions for all relevant robustness check procedures. In addition, editorial boards and reviewers should consider adopting a more conservative stance when it comes to reviewing PLS-SEM studies, ensuring that appropriate robustness checks are performed where applicable.

This study has certain limitations. First, the journals included in our SLR may not be representative of all journals that published PLS-SEM studies within the investigated timeframe and field of study. Owing to the impracticality of considering all such journals, we considered only selected high-ranking publications.

Second, even though we conducted our SLR search within Clarivate Analytics' Web of Science Core Collection, we did the selection of high-ranking publications, using Elsevier's SCImago. Nevertheless, by doing so, we have broadened the range of journals that were taken into account, as SCImago is less conservative than Clarivate's Journal Citation Reports (JCR) when it comes to ranking journals in the top-tier zone.

Third, even though some scholars (e.g., Hair et al., 2020) suggest that robustness checks should also include multicollinearity, by examining bivariate correlations between formative indicators or between exogenous constructs, multicollinearity was not part of our analysis, as its assessment is generally associated to PLS structural and measurement model assessment. Moreover, the necessity of bivariate correlations between independent variables in a regression model as a condition for multicollinearity is a subject of debate (e.g., Gujarati & Porter, 2009). Nevertheless, further research and refinement of techniques for examining multicollinearity in PLS-SEM models would be beneficial.

Finally, the robustness checks covered in this paper are not meant to be exhaustive; rather, they were selected based on their relevance within the conventional regression analysis framework. When using PLS-SEM, researchers should also consider other relevant robustness checks. For instance, they should check for issues related to common method variance (Chin et al., 2013), assess whether any independent variable in the model is necessary to produce a specific outcome for a dependent variable (necessary condition analysis - NCA; Richter et al., 2020), and examine whether the specification (i.e., formative/reflective) of the measurement model is correct (confirmatory tetrad analysis - CTA; Gudergan et al., 2008). Additionally, researchers should assess whether their model, when compared with other competing models, performs better in terms of information criteria (Sharma et al., 2019), or predictive ability (PLSpredict; Shmueli et al., 2019; cross validated predictive ability test - CVPAT; Liengard et al., 2021). Future research exploring

**Table 5**  
Number of instances heteroskedasticity checks were performed.

Technique Used	2016–2018		2019–2021	
	Count	%	Count	%
Levene	1	33	5	45
Graphical method	1	33	3	27
Details not provided	0	0	2	18
Breusch-Pagan	1	33	0	0
Koenker	0	0	1	9
<b>Total</b>	<b>3</b>	<b>100</b>	<b>11</b>	<b>100</b>

**Table 6**  
Robustness checks recommendations.

When to Test	Recommended technique(s)	Tool(s)	Additional technique(s)	Tool(s)
<b>Normality – univariate</b>				
At all times	Examination of skewness and kurtosis values	✓ PLS-SEM programs (e.g., SmartPLS) ✓ Microsoft Excel functions - SKEW() and KURT() ✓ <i>skewness()</i> and <i>kurtosis()</i> in R 'moments' package	Shapiro-Wilk Kolmogorov-Smirnov	✓ <i>shapiro.test()</i> in R 'stats' package ✓ <i>ks.test()</i> in R 'stats' package
<b>Normality – multivariate</b>				
Univariate normality failed to be rejected	Mardia's multivariate normality test	✓ Webpower website <a href="https://webpower.psychstat.org/models/kurtosis/">https://webpower.psychstat.org/models/kurtosis/</a> ✓ <i>mvnTest()</i> in R 'MVN' package	Small (1980), Srivastava (1984)	R codes not available in any existing packages
<b>Endogeneity</b>				
Objective of study is confirmatory rather than predictive	Gaussian copula approach (if assumptions of the approach are met)	✓ PLS-SEM programs (e.g., SmartPLS) ✓ R codes are available at <a href="https://www.pls-sem.net/downloads/gaussian-copula-files/">https://www.pls-sem.net/downloads/gaussian-copula-files/</a>	Durbin-Wu-Hausman Test	✓ Using the generic <i>summary()</i> where the object is an instrumental variable regression estimated using R 'ivreg' package
<b>Non-linearity</b>				
At all times	RESET test followed by Higher order polynomial fitting (quadratic)	✓ <i>resettest()</i> in R 'zoo' package ✓ PLS-SEM programs (e.g., SmartPLS) – Assess the significance of quadratic terms	–	–
<b>Unobserved heterogeneity</b>				
At all times	FIMIX-PLS followed by PLS-POS	✓ PLS-SEM programs (e.g., SmartPLS) – Detect unobserved heterogeneity and number of segments ✓ PLS-SEM programs (e.g., SmartPLS) – Determine the actual structure of the segments	– REBUS-PLS PLS-GASPLS-IRRS	– ✓ <i>rebus.pls()</i> in R 'plspm' package R codes not available in any existing packages

the use of such robustness checks in business-related studies would be beneficial for enhancing PLS-SEM rigor in business research.

**Funding**

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

**CRedit authorship contribution statement**

**Santha Vaithilingam:** Writing – review & editing, Writing – original draft, Supervision, Methodology, Conceptualization. **Chu Sun Ong:** Writing – original draft, Writing - review & editing, Methodology, Formal Analysis. **Ovidiu I. Moisescu:** Writing – review & editing, Writing – original draft, Conceptualization. **Mahendhiran S. Nair:** Writing – review & editing, Supervision, Methodology.

**Declaration of Competing Interest**

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

**Acknowledgement**

The authors would like to extend their gratitude to our research assistants Ms. Chin Si Yen and Mr. Kaveendra Vasuthevan for their excellent support provided to the second author in the systematic review process including locating, coding, retrieving metadata, and identifying the robustness checks performed and associated techniques used in the sample studies. The work of Ovidiu I. Moisescu was supported by a grant of the Ministry of Research, Innovation and Digitization, CNCS - UEFISCDI, project number PN-III-P1-1.1-TE-2021-0271, within PNCDI III.

**Appendix A. Robustness checks in PLS-SEM empirical studies published between 2016 and 2021 in business journals**

	Number of PLS-SEM empirical papers	Number of papers performing at least one robustness check	Number of instances robustness checks were performed
<i>Journal of Business Research</i>	139	23	32
<i>Technological Forecasting and Social Change</i>	51	12	21
<i>Journal of Retailing and Consumer Services</i>	108	11	18
<i>Journal of Entrepreneurship in Emerging Economies</i>	29	11	13
<i>European Journal of Marketing</i>	19	8	11
<i>Journal of Business &amp; Industrial Marketing</i>	60	7	8
<i>Journal of Product and Brand Management</i>	39	6	7
<i>Business Strategy and the Environment</i>	29	6	7
<i>Internet Research</i>	50	4	6
<i>Journal of Services Marketing</i>	36	6	6

(continued on next page)

(continued)

	Number of PLS-SEM empirical papers	Number of papers performing at least one robustness check	Number of instances robustness checks were performed
<i>Journal of Intellectual Capital</i>	38	4	5
<i>Corporate Social Responsibility and Env. Management</i>	34	4	5
<i>Industrial Marketing Management</i>	23	5	5
<i>European Business Review</i>	22	3	5
<i>Journal of Business Ethics</i>	17	4	5
<i>South Asian Journal of Business Studies</i>	11	3	5
<i>International Journal of Accounting Information Systems</i>	7	2	5
<i>International Journal of Retail &amp; Distribution Management</i>	37	3	4
<i>Journal of Service Theory and Practice</i>	14	3	4
<i>Journal of Small Business and Enterprise Development</i>	11	2	4
<i>EuroMed Journal of Business</i>	10	3	4
<i>Journal of Hospitality Marketing &amp; Management</i>	9	3	4
<i>Long Range Planning</i>	8	2	4
<i>Thunderbird International Business Review</i>	8	2	4
<i>Research in Transportation Business and Management</i>	6	3	4
<i>Small Business Economics</i>	4	2	4
<i>Journal of Vacation Marketing</i>	17	3	3
<i>International Journal of Managing Projects in Business</i>	14	3	3
<i>Electronic Commerce Research and Applications</i>	13	1	3
<i>International Business Review</i>	9	3	3
<i>Management Decision</i>	44	2	2
<i>European Management Journal</i>	21	2	2
<i>Journal of Asia Business Studies</i>	19	2	2
<i>Journal of Marketing Management</i>	13	1	2
<i>Journal of Electronic Commerce Research</i>	11	1	2
<i>Business Strategy and Development</i>	9	1	2
<i>IEEE Transactions on Engineering Management</i>	7	2	2
<i>British Journal of Management</i>	6	2	2
<i>Journal of Engineering and Technology Management</i>	6	2	2
<i>R &amp; D Management</i>	4	2	2
<i>Business Process Management Journal</i>	31	1	1
<i>International Entrepreneurship and Management Journal</i>	24	1	1
<i>Young Consumers</i>	16	1	1
<i>Int. J. of Entrepreneurial Behavior &amp; Research</i>	14	1	1
<i>Electronic Markets</i>	9	1	1
<i>Journal of Fashion Marketing and Management</i>	8	1	1
<i>Journal of World Business</i>	5	1	1
<i>Journal of Entrepreneurship</i>	3	1	1
<i>Emerging Markets Finance and Trade</i>	2	1	1
<i>International Journal of Advertising</i>	2	1	1
<i>Journal of International Business Studies</i>	2	1	1
<i>Journal of the Academy of Marketing Science</i>	2	1	1
<i>Asian Business &amp; Management</i>	1	1	1
<i>European Research on Management and Bus. Economics</i>	12	0	0
<i>International Journal of Management Education</i>	11	0	0
<i>Psychology &amp; Marketing</i>	10	0	0
<i>International Marketing Review</i>	9	0	0
<i>Journal of International Entrepreneurship</i>	9	0	0
<i>Journal of African Business</i>	8	0	0
<i>Service Business</i>	7	0	0
<i>BRQ-Business Research Quarterly</i>	6	0	0
<i>Eurasian Business Review</i>	4	0	0
<i>Journal of Innovation &amp; Knowledge</i>	3	0	0
<i>International Small Business Journal</i>	2	0	0
<i>Journal of Global Fashion Marketing</i>	2	0	0
<i>Journal of Interactive Marketing</i>	2	0	0
<i>Journal of Media Business Studies</i>	2	0	0
<i>Multinational Business Review</i>	2	0	0
<i>Journal of Business and Psychology</i>	1	0	0
<i>Journal of Family Business Strategy</i>	1	0	0
<i>Journal of Management Analytics</i>	1	0	0
<i>Journal of Marketing</i>	1	0	0
<i>Journal of Personal Selling &amp; Sales Management</i>	1	0	0
<i>Journal of Product Innovation Management</i>	1	0	0
<i>Journal of Service Research</i>	1	0	0
<i>Public Relations Review</i>	1	0	0
<b>Total</b>	<b>1,228</b>	<b>182</b>	<b>245</b>

## References

- Ali, F., Rasoolimanesh, S. M., Sarstedt, M., Ringle, C. M., & Ryu, K. (2018). An assessment of the use of partial least squares structural equation modeling (PLS-SEM) in hospitality research. *International Journal of Contemporary Hospitality Management*, 30(1), 514–538. <https://doi.org/10.1108/IJCHM-10-2016-0568>
- Basco, R., Hair, J. F., Ringle, C. M., & Sarstedt, M. (2022). Advancing family business research through modeling nonlinear relationships: Comparing PLS-SEM and multiple regression. *Journal of Family Business Strategy*, 13(3), Article 100457. <https://doi.org/10.1016/j.jfbs.2021.100457>
- Bayonne, E., Marin-García, J. A., & Alfalla-Luque, R. (2020). Partial least squares (PLS) in Operations Management research: Insights from a systematic literature review. *Journal of Industrial Engineering and Management*, 13(3), 565–597. <https://doi.org/10.3926/jiem.3416>
- Becker, J. M., Proksch, D., & Ringle, C. M. (2022). Revisiting Gaussian copulas to handle endogenous regressors. *Journal of the Academy of Marketing Science*, 50, 46–66. <https://doi.org/10.1007/s11747-021-00805-y>
- Becker, J.-M., Rai, A., Ringle, C. M., & Völckner, F. (2013). Discovering Unobserved Heterogeneity in Structural Equation Models to Avert Validity Threats. *MIS Quarterly*, 37(3), 665–694. <https://doi.org/10.25300/MISQ/2013/37.3.01>
- Cepeda-Carrión, G., Cegarra-Navarro, J. G., & Cillo, V. (2018). Tips to use partial least squares structural equation modelling (PLS-SEM) in knowledge management. *Journal of Knowledge Management*, 23(1), 67–89. <https://doi.org/10.1108/JKM-05-2018-0322>
- Cepeda-Carrión, G., Henseler, J., Ringle, C. M., & Roldán, J. L. (2016). Prediction-oriented modeling in business research by means of PLS path modeling: Introduction to a JBR special section. *Journal of Business Research*, 69(10), 4545–4551. <https://doi.org/10.1016/j.jbusres.2016.03.048>
- Certo, S. T., Busenbark, J. R., Woo, H.-S., & Semadeni, M. (2016). Sample selection bias and Heckman models in strategic management research. *Strategic Management Journal*, 37(13), 2639–2657. <https://doi.org/10.1002/smj.2475>
- Chin, W. W., Thatcher, J. B., Wright, R. T., & Steel, D. (2013). Controlling for common method variance in PLS analysis: The measured latent marker variable approach. In H. Abdi, W. W. Chin, V. Esposito Vinzi, G. Russolillo, & L. Trinchera (Eds.), *New perspectives in partial least squares and related methods* (pp. 231–239). Springer. [https://doi.org/10.1007/978-1-4614-8283-3\\_16](https://doi.org/10.1007/978-1-4614-8283-3_16)
- DeCarlo, L. T. (1997). On the meaning and use of kurtosis. *Psychological Methods*, 2(3), 292–307. <https://doi.org/10.1037/1082-989x.2.3.292>
- Dessart, L., Aldas-Manzano, J., & Veloutsou, C. (2019). Unveiling heterogeneous engagement-based loyalty in brand communities. *European Journal of Marketing*, 53(9), 1854–1881. <https://doi.org/10.1108/EJM-11-2017-0818>
- Ebbes, P., Papies, D., & van Heerde, H. J. (2016). Dealing with Endogeneity: A nontechnical Guide for Marketing Researchers. In C. Homburg, M. Klarmann, & A. Vomberg (Eds.), *Handbook of Market Research* (pp. 1–37). Springer. [https://doi.org/10.1007/978-3-319-05542-8\\_8-1](https://doi.org/10.1007/978-3-319-05542-8_8-1)
- Ebbes, P., Papies, D., & van Heerde, H. J. (2011). The Sense and Non-Sense of Holdout Sample Validation in the Presence of Endogeneity. *Marketing Science*, 30(6), 1115–1122. <https://doi.org/10.1287/mksc.1110.0666>
- Esposito Vinzi, V., Trinchera, L., Squillacioti, S., & Tenenhaus, M. (2008). REBUS-PLS: A response-based procedure for detecting unit segments in PLS path modelling. *Applied Stochastic Models in Business and Industry*, 24(5), 439–458. <https://doi.org/10.1002/asmb.728>
- Gudergan, S. P., Ringle, C. M., Wende, S., & Will, A. (2008). Confirmatory Tetrad Analysis in PLS Path Modeling. *Journal of Business Research*, 61(12), 1238–1249. <https://doi.org/10.1016/j.jbusres.2008.01.012>
- Guenther, P., Guenther, M., Ringle, C. M., Zaeferian, G., & Cartwright, S. (2023). Improving PLS-SEM use for business marketing research. *Industrial Marketing Management*, 111, 127–142. <https://doi.org/10.1016/j.indmarman.2023.03.010>
- Guide, V. D. R., & Ketokivi, M. (2015). Notes from the Editors: Redefining some methodological criteria for the journal. *Journal of Operations Management*, 37(1), v–viii. [https://doi.org/10.1016/S0272-6963\(15\)00056-X](https://doi.org/10.1016/S0272-6963(15)00056-X)
- Gujarati, D. N., & Porter, D. C. (2009). *Basic Econometrics* (5th ed.). McGraw-Hill Irwin.
- Hair, J. F., Astrachan, C. B., Moisescu, O. I., Radomir, L., Sarstedt, M., Vaithilingam, S., & Ringle, C. M. (2021). Executing and interpreting applications of PLS-SEM: Updates for family business researchers. *Journal of Family Business Strategy*, 12(3), Article 100392. <https://doi.org/10.1016/j.jfbs.2020.100392>
- Hair, J. F., Babin, B. J., Anderson, R. E., & Black, W. C. (2019a). *Multivariate Data Analysis* (8th ed.). Cengage Learning.
- Hair, J. F., Hult, T., Ringle, C. M., & Sarstedt, M. (2022). *A Primer on Partial Least Squares Structural Equation Modeling (PLS-SEM)* (3rd ed.). Thousand Oaks: Sage.
- Hair, J. F., Risher, J. J., Sarstedt, M., & Ringle, C. M. (2019b). When to use and how to report the results of PLS-SEM. *European Business Review*, 31(1), 2–24. <https://doi.org/10.1108/EBR-11-2018-0203>
- Hair, J. F., Sarstedt, M., & Ringle, C. M. (2019c). Rethinking some of the rethinking of partial least squares. *European Journal of Marketing*, 53(4), 566–584. <https://doi.org/10.1108/Ejm-10-2018-0665>
- Hair, J. F., Sarstedt, M., Matthews, L. M., & Ringle, C. M. (2016). Identifying and treating unobserved heterogeneity with FIMIX-PLS: Part I—method. *European Business Review*, 28(1), 63–76. <https://doi.org/10.1108/EBR-09-2015-0094>
- Hair, J. F., Sarstedt, M., Pieper, T. M., & Ringle, C. M. (2012). The use of partial least squares structural equation modeling in strategic management research: A review of past practices and recommendations for future applications. *Long Range Planning*, 45(5–6), 320–340. <https://doi.org/10.1016/j.lrp.2012.09.008>
- Hair, J. F., Sarstedt, M., Ringle, C. M., & Gudergan, S. P. U. (2024). *Advanced Issues in Partial Least Squares Structural Equation Modeling (PLS-SEM)* (2nd ed.). SAGE Publications.
- Hair, J. F., Hollingsworth, C. L., Randolph, A. B., & Chong, A. Y. L. (2017). An updated and expanded assessment of PLS-SEM in information systems research. *Industrial Management + Data Systems*, 117(3), 442–458. <https://doi.org/10.1108/IMDS-04-2016-0130>
- Hair, J. F., Howard, M. C., & Nitzl, C. (2020). Assessing measurement model quality in PLS-SEM using confirmatory composite analysis. *Journal of Business Research*, 109, 101–110. <https://doi.org/10.1016/j.jbusres.2019.11.069>
- Henseler, J. (2018). Partial least squares path modeling: Quo vadis? *Quality & Quantity*, 52(1), 1–8. <https://doi.org/10.1007/s11135-018-0689-6>
- Hult, G. T. M., Hair, J. F., Proksch, D., Sarstedt, M., Pinkwart, A., & Ringle, C. M. (2018). Addressing Endogeneity in International Marketing Applications of Partial Least Squares Structural Equation Modeling. *Journal of International Marketing*, 26(3), 1–21. <https://doi.org/10.1509/jim.17.0151>
- Kaufmann, L., & Gaeckler, J. (2015). A structured review of partial least squares in supply chain management research. *Journal of Purchasing and Supply Management*, 21(4), 259–272. <https://doi.org/10.1016/j.pursup.2015.04.005>
- Kitchenham, B., Brereton, O. P., Budgen, D., Turner, M., Bailey, J., & Linkman, S. (2009). Systematic literature reviews in software engineering - A systematic literature review. *Information and Software Technology*, 51(1), 7–15. <https://doi.org/10.1016/j.infsof.2008.09.009>
- Kock, N. (2015). *WarpPLS 5.0 User Manual*. Laredo, TX: ScriptWarp Systems.
- Kock, N. (2022). *WarpPLS User Manual: Version 8.0*. Laredo, TX: ScriptWarp Systems.
- Latan, H. (2018). PLS Path Modeling in Hospitality and Tourism Research: The Golden Age and Days of Future Past. In F. Ali, S. M. Rasoolimanesh, & C. Cobanoglu (Eds.), *Applying Partial Least Squares in Tourism and Hospitality Research* (pp. 53–83). Emerald Publishing Limited. <https://doi.org/10.1108/978-1-78756-699-620181004>
- Liengaard, B. D., Sharma, P. N., Hult, G. T. M., Jensen, M. B., Sarstedt, M., Hair, J. F., & Ringle, C. M. (2021). Prediction: Coveted, Yet Forsaken? Introducing a Cross-validated Predictive Ability Test in Partial Least Squares Path Modeling. *Decision Sciences*, 52(2), 362–392. <https://doi.org/10.1111/deci.12445>
- Loy, A., Follett, L., & Hofmann, H. (2016). Variations of Q-Q Plots: The Power of Our Eyes! *American Statistician*, 70(2), 202–214. <https://doi.org/10.1080/00031305.2015.1077728>
- Mardia, K. V. (1970). Measures of Multivariate Skewness and Kurtosis with Applications. *Biometrika*, 57(3), 519–530. <https://doi.org/10.2307/2334770>
- Mardia, K. V. (1974). Applications of Some Measures of Multivariate Skewness and Kurtosis in Testing Normality and Robustness Studies. *Sankhya—the Indian Journal of Statistics Series B (1960–2002)*, 36(2), 115–128. <http://www.jstor.org/stable/25051892>
- Matthews, L. M., Sarstedt, M., Hair, J. F., & Ringle, C. M. (2016). Identifying and treating unobserved heterogeneity with FIMIX-PLS: Part II—A case study. *European Business Review*, 28(2), 63–76. <https://doi.org/10.1108/EBR-09-2015-0095>
- Mikalef, P., & Pateli, A. (2017). Information technology-enabled dynamic capabilities and their indirect effect on competitive performance: Findings from PLS-SEM and fsQCA. *Journal of Business Research*, 70, 1–16. <https://doi.org/10.1016/j.jbusres.2016.09.004>
- Papies, D., Ebbes, P., & Van Heerde, H. J. (2017). Addressing Endogeneity in Marketing Models. In P. S. H. Leeflang, J. E. Wieringa, T. H. A. Bijmolt, & K. H. Pauwels (Eds.), *Advanced Methods for Modeling Markets* (pp. 581–627). Springer International Publishing. [https://doi.org/10.1007/978-3-319-53469-5\\_18](https://doi.org/10.1007/978-3-319-53469-5_18)
- Park, S., & Gupta, S. (2012). Handling Endogenous Regressors by Joint Estimation Using Copulas. *Marketing Science*, 31(4), 567–586. <https://doi.org/10.1287/mksc.1120.0718>
- Pierce, J. R., & Aguinis, H. (2013). The Too-Much-of-a-Good-Thing Effect in Management. *Journal of Management*, 39(2), 313–338. <https://doi.org/10.1177/0149206311410060>
- Ramsey, J. B. (1969). Tests for Specification Errors in Classical Linear Least-Squares Regression Analysis. *Journal of the Royal Statistical Society. Series B (Methodological)*, 31(2), 350–371. <http://www.jstor.org/stable/2984219>
- Razali, N. M., & Wah, Y. B. (2011). Power Comparisons of Shapiro-Wilk, Kolmogorov-Smirnov, Lilliefors and Anderson-Darling tests. *Journal of Statistical Modeling and Analytics*, 2(1), 21–33.
- Richter, N. F., Schubring, S., Hauff, S., Ringle, C. M., & Sarstedt, M. (2020). When Predictors of Outcomes are Necessary: Guidelines for the Combined use of PLS-SEM and NCA. *Industrial Management + Data Systems*, 120(12), 2243–2267. <https://doi.org/10.1108/IMDS-11-2019-0638>
- Richter, N. F., Hauff, S., Ringle, C. M., & Gudergan, S. P. (2022). The Use of Partial Least Squares Structural Equation Modeling and Complementary Methods in International Management Research. *Management International Review*, 62, 449–470. <https://doi.org/10.1007/s11575-022-00475-0>
- Ringle, C. M., Sarstedt, M., & Mooi, E. A. (2010). Response-Based Segmentation Using Finite Mixture Partial Least Squares: Theoretical Foundations and an Application to American Customer Satisfaction Index Data. In R. Stahlbock, S. Crone, & S. Lessmann (Eds.), *Data Mining. Annals of Information Systems* (pp. 19–49). Springer. [https://doi.org/10.1007/978-1-4419-1280-0\\_2](https://doi.org/10.1007/978-1-4419-1280-0_2)

- Ringle, C. M., Sarstedt, M., & Schlittgen, R. (2014). Genetic algorithm segmentation in partial least squares structural equation modeling. *OR spectrum*, 36(1), 251–276. <https://doi.org/10.1007/s00291-013-0320-0>
- Ringle, C. M., Sarstedt, M., Mitchell, R., & Gudergan, S. P. (2020). Partial least squares structural equation modeling in HRM research. *International Journal of Human Resource Management*, 31(12), 1617–1643. <https://doi.org/10.1080/09585192.2017.1416655>
- Ringle, C. M., Sarstedt, M., Sinkovics, N., & Sinkovics, R. R. (2023). A perspective on using partial least squares structural equation modelling in data articles. *Data in Brief*, 48, Article 109074. <https://doi.org/10.1016/j.dib.2023.109074>
- Ringle, C. M., Wende, S. & Becker, J. M. (2022). *SmartPLS* (Version 4) [Computer Software] Oststeinbek: SmartPLS. Retrieved from <https://www.smartpls.com>.
- Sarstedt, M., & Cheah, J. H. (2019). Partial least squares structural equation modeling using SmartPLS: a software review. *Journal of Marketing Analytics*, 7, 196–202. <https://doi.org/10.1057/s41270-019-00058-3>
- Sarstedt, M., & Mooi, E. (2014). *A Concise Guide to Market Research: The Process, Data, and Methods Using IBM SPSS Statistics*. Berlin Heidelberg: Springer.
- Sarstedt, M., Becker, J.-M., Ringle, C. M., & Schwaiger, M. (2011). Uncovering and treating unobserved heterogeneity with FIMIX-PLS: Which model selection criterion provides an appropriate number of segments? *Schmalenbach Business Review*, 63(1), 34–62. <https://doi.org/10.1007/BF03396886>
- Sarstedt, M., Hair, J. F., Pick, M., Liengaard, B. D., Radomir, L., & Ringle, C. M. (2022a). Progress in partial least squares structural equation modeling use in marketing research in the last decade. *Psychology & Marketing*, 39(5), 1035–1064. <https://doi.org/10.1002/mar.21640>
- Sarstedt, M., Radomir, L., Moisescu, O. I., & Ringle, C. M. (2022b). Latent class analysis in PLS-SEM: A review and recommendations for future applications. *Journal of Business Research*, 138, 398–407. <https://doi.org/10.1016/j.jbusres.2021.08.051>
- Sarstedt, M., Ringle, C. M., Cheah, J. H., Ting, H. R., Moisescu, O. I., & Radomir, L. (2020). Structural model robustness checks in PLS-SEM. *Tourism Economics*, 26(4), 531–554. <https://doi.org/10.1177/1354816618823921>
- Schlittgen, R., Ringle, C. M., Sarstedt, M., & Becker, J.-M. (2016). Segmentation of PLS path models by iterative reweighted regressions. *Journal of Business Research*, 69(10), 4583–4592. <https://doi.org/10.1016/j.jbusres.2016.04.009>
- SCImago (n.d.). *SJR — SCImago Journal & Country Rank* [Portal]. Retrieved January 29, 2022, from <http://www.scimagojr.com>.
- Schubert, F., Rademaker, M. E., & Henseler, J. (2022). Assessing the overall fit of composite models estimated by partial least squares path modeling. *European Journal of Marketing*, 57(6), 46–66. <https://doi.org/10.1108/ejm-08-2020-0586>
- Sharma, P. N., Sarstedt, M., Shmueli, G., Kim, K. H., & Thiele, K. O. (2019). PLS-Based Model Selection: The Role of Alternative Explanations in Information Systems Research. *Journal of the Association for Information Systems*, 20(4), 346–397. <https://doi.org/10.17005/1.jais.00538>
- Shmueli, G. (2010). To Explain or to Predict? *Statistical Science*, 25(3), 289–310. <https://doi.org/10.1214/10-Sts330>
- Shmueli, G., Ray, S., Estrada, J. M. V., & Chatla, S. B. (2016). The elephant in the room: Predictive performance of PLS models. *Journal of Business Research*, 69(10), 4552–4564. <https://doi.org/10.1016/j.jbusres.2016.03.049>
- Shmueli, G., Sarstedt, M., Hair, J. F., Cheah, J. H., Ting, H., Vaithilingam, S., & Ringle, C. M. (2019). Predictive Model Assessment in PLS-SEM: Guidelines for Using PLSpredict. *European Journal of Marketing*, 53(11), 2322–2347. <https://doi.org/10.1108/EJM-02-2019-0189>
- Small, N. J. H. (1980). Marginal Skewness and Kurtosis in Testing Multivariate Normality. *Journal of the Royal Statistical Society. Series C (Applied Statistics)*, 29(1), 85–87. <https://doi.org/10.2307/2346414>
- Srivastava, M. S. (1984). A Measure of Skewness and Kurtosis and a Graphical-Method for Assessing Multivariate Normality. *Statistics & Probability Letters*, 2(5), 263–267. [https://doi.org/10.1016/0167-7152\(84\)90062-2](https://doi.org/10.1016/0167-7152(84)90062-2)
- Svensson, G., Ferro, C., Høgevoid, N., Padin, C., Varela, J. C. S., & Sarstedt, M. (2018). Framing the triple bottom line approach: Direct and mediation effects between economic, social and environmental elements. *Journal of Cleaner Production*, 197, 972–991. <https://doi.org/10.1016/j.jclepro.2018.06.226>
- Vaithilingam, S., Nair, M., Macharia, M., & Venkatesh, V. (2022). Mobile communication and use behavior of the urban poor in a developing country: A field study in Malaysia. *International Journal of Information Management*, 63, Article 102440. <https://doi.org/10.1016/j.ijinfomgt.2021.102440>
- West, R. E., & Rich, P. J. (2012). Rigor, Impact and Prestige: A Proposed Framework for Evaluating Scholarly Publications. *Innovative Higher Education*, 37, 359–371. <https://doi.org/10.1007/s10755-012-9214-3>
- Wooldridge, J. M. (2010). *Econometric Analysis of Cross Section and Panel Data* (2nd ed.). MIT Press.
- Wooldridge, J. M. (2013). *Introductory Econometrics: A Modern Approach* (5th ed.). Cengage Learning.
- Santha Vaithilingam** is Professor of Econometrics and Policy Modelling, at Sunway Institute for Global Strategy and Competitiveness, Sunway University. She is trained in mathematical modeling. Her current research interest revolves around behavioral econometrics, using mathematical and statistical modeling to understand human behavior and the decision-making process. Her research work is also focused on developing research instruments and using robust data analytics & advanced statistical tools to research technology, innovation, and economic policy modeling for enhancing sustainable economic development.
- Chu Sun Ong** is a Ph.D. student at the Tokyo Institute of Technology. His doctoral research focuses on modelling the societal implications of digital inequality. He holds a bachelor's degree majoring in Econometrics and Business Statistics and a master's degree in Economics.
- Ovidiu I. Moisescu** is Professor of Marketing and Branding at the Marketing Department, Faculty of Economics and Business Administration at Babeş-Bolyai University of Cluj-Napoca, Romania. He completed his PhD in Marketing at the West University of Timișoara (Romania), as well as a postdoctoral research project at Babeş-Bolyai University of Cluj-Napoca (Romania) and Corvinus University of Budapest (Hungary). His research focuses on consumer behavior, brand loyalty, CSR, and methodological issues in PLS-SEM.
- Mahendhiran S. Nair** is Professor and Pro Vice-Chancellor (Research Engagement and Impact) at Sunway University. He is trained as an econometrician and data scientist. His expertise and research interests include data science; econometrics; behavioral economics; foresighting, science & technology policy modeling; planetary health; and sustainable development. He has been a subject matter expert for government agencies, public policy organizations and 'think-tanks' in Malaysia and regional economies. He has led to the formulation of several national economic & technology plans and studies for Malaysia.